

African Gender Studies Quarterly



Editorial Note

It is with great joy that we introduce this inaugural edition of African Gender Studies Quarterly (AGenSQ), a journal established to provide a dedicated platform for rigorous, interdisciplinary scholarship on gender and its intersections with African societies and the broader Global South. As a quarterly publication, we are committed to fostering critical dialogue and advancing knowledge that is both theoretically grounded and contextually relevant.

In this first issue, we are pleased to present a collection of articles that exemplify the depth and diversity of gender scholarship. The opening paper, "A Biopsychosocial Perspective on Sexual Disorders and Dysfunctions: Implications for Diagnosis and Intervention," offers a comprehensive examination of sexual health through biological, psychological, and sociocultural lenses, emphasizing the need for culturally informed care in African contexts. The authors make a compelling case for destigmatising sexual health discussions and integrating multidisciplinary approaches to treatment.

The second article, "Love in the Digital Age: Rethinking Courtship and Romance in Contemporary Society," explores how technology, shifting gender roles, and evolving social norms are transforming intimate relationships. By examining phenomena such as digital dating, emotional labour, and self-love, this paper illuminates the complex terrain of modern romance and its implications for psychological well-being.

Following this, "Cultural Influences on the Perception, Regulation, and Communication of Sexual Content" investigates the powerful role of media in shaping sexual attitudes and behaviours. Drawing on psychological theories, the authors critically analyse how different media forms from social media to pornography influence sexual socialisation, particularly among young people, and advocate for enhanced media literacy and psychological interventions.

The fourth contribution, "Marriage, Sexuality, and Cultural Norms in Indigenous Nigerian Films: A Comparative Analysis of Yoruba, Hausa, and Igbo Cinema," employs qualitative content analysis to examine how indigenous films reflect and reinforce cultural values surrounding marriage and sexuality. The study reveals persistent gender double standards while noting emergent progressive shifts, demonstrating the pedagogical function of cinema in sustaining or challenging societal norms.

The fifth article, "Gendered Constructed Strategies in Conflict Resolution: Challenges Faced by Female Secondary School Leaders in Chivi, Zimbabwe," illuminates how gender shapes leadership in educational settings. Through rich qualitative data, the authors identify three distinct gendered strategies employed by female school heads: the motherly care approach, the relational strategy, and the inclusive gender-sensitive approach.

The sixth article, "Political Education and Voters' Behaviour in Nigeria," establishes a strong positive relationship between political education and voter behaviour. Notably,

with women constituting 64.5% of respondents, the findings invite critical reflection on gendered patterns of political participation and the need for gender-sensitive civic education approaches.

We believe these articles have been thoroughly reviewed, and the recommendations offered will help to properly formulate effective solutions to identified issues. This inaugural issue has done well by providing a space where both emerging and established scholars alike have engaged to present themes that demand attention and open engagement. We sincerely hope that this collection will not only inform but also invite deeper reflection and continued scholarship on the evolving intersections of gender and its dynamics as they affect Africa and the rest of the third world.

Dr. O. A.T.Olatunji,
Editor, African Gender Studies Quarterly,
Associate Professor / Reader @ Department of Psychology,
Faculty of the Social Sciences,
Federal University, Oye Ekiti, Ekiti State,
Nigeria.
Email: ajoke.olatunji@fuoye.edu.ngvii

Table of Content

1. Biopsychosocial Perspective on Sexual Disorders and Dysfunctions: Implications for Diagnosis and Intervention Wosangha S. Otu, Idoko R. Ihotu & Okonkwo K. Ifunanya	5
2. Love in the Digital Age: Rethinking Courtship and Romance in Contemporary Society Olaleye Fuad Duncan, Adekanmbi Adebawale Isaac & Lijoka Tomilola Joshua	19
3. Cultural Influences on the Perception, Regulation, and Communication of Sexual Content OBE, O., EKOTT, D. & Adegbohun, O. J.	38
4. Marriage, Sexuality, and Cultural Norms in Indigenous Nigerian Films: A Comparative Analysis of Yoruba, Hausa, and Igbo Cinema Stephen, Emmanuel Ayomide & Osaghae, Japheth Osasere	52
5. Gendered constructed strategies in conflict resolution: Challenges faced by female secondary school leaders in Chivi, Zimbabwe Clemence Dzingirai, Rosemary Guvhu, Anna Moyo,	66
6. Political Education and Voters' Behaviour in Nigeria: A Case Study of the 2019 General Election in Lagos State Deborah Ifeoluwa, Makanjuola	80

DOI: <https://doi.org/10.69778/3106-7603/2026/2.1/a1>

BIOPSYCHOSOCIAL PERSPECTIVE ON SEXUAL DISORDERS AND DYSFUNCTIONS: IMPLICATIONS FOR DIAGNOSIS AND INTERVENTION

Wosangha S. Otu,
Department of Psychology,
University of Lagos
Email: wosangha16@gmail.com

Idoko R. Ihotu
Department of Psychology,
University of Lagos

Okonkwo K. Ifunanya
Department of Psychology,
University of Lagos

Abstract

Sexual disorders and dysfunctions are conditions that disrupt satisfaction in sexual activity and often affect self-esteem, intimate relationships, and overall quality of life. These disorders encompass difficulties related to sexual desire, arousal, orgasm, pain during intercourse, and certain paraphilic behaviors. Although often underreported due to stigma and cultural sensitivities, sexual dysfunction represents a significant public health concern affecting individuals across different age groups and sociocultural contexts. The aetiology of these conditions is multifaceted, involving interactions among biological, psychological, and sociocultural factors. Biological contributors may include hormonal imbalances, neurological or vascular impairments, chronic health conditions, and medication side effects. Psychological influences such as performance anxiety, past traumatic experiences, negative body image, depression, and low self-esteem may further exacerbate sexual difficulties. Additionally, sociocultural factors, including relationship conflicts, rigid gender expectations, religious beliefs, and societal stigma surrounding sexuality often shape individuals' attitudes toward sexual expression and discourage help-seeking behaviour. Accurate diagnosis requires a comprehensive, culturally sensitive clinical assessment that integrates medical and psychosocial perspectives. Standardized evaluation tools such as the Female Sexual Function Index (FSFI) and the International Index of Erectile Function (IIEF) are often employed to support systematic assessment and guide clinical decision-making. Treatment approaches are typically multidimensional, combining psychological interventions, medical management, physiotherapy, and, increasingly, digital health technologies that expand access to care and education. This paper underscores the importance of adopting a biopsychosocial framework in understanding and managing sexual disorders and dysfunctions. By emphasising culturally informed care, reducing stigma, and promoting open discussions about sexual health, such an approach can enhance therapeutic outcomes and contribute to improved psychological well-being, relational satisfaction, and overall quality of life.

Keywords: *Sexual disorders, sexual dysfunction, biopsychosocial model, sexual health, cultural factors,*

INTRODUCTION

Sexuality, personal and intimate, is of the utmost importance to the makeup of human beings. However, for many, sexual functioning can become a source of genuine hurt and frustration, thereby making what we call sexual disorders and dysfunctions. These are significant public health issues affecting many people of all genders and ages worldwide. By such problems, we allude to the persistent or recurrent disturbances in sexual desire and arousal or in the capacity to attain orgasm or to pass through ejaculation or even experiencing pain during intercourse. These are not merely physical symptoms but also significantly affect one's quality of life, self-esteem, and intimate relationships (American Psychiatric Association, 2013). Hence, understanding sexual disorders goes beyond the clinical and therapeutic stance because they impinge on reproduction, relationship stability, and mental health.

Quite obviously, sexual difficulties touch a great population and hugely affect the quality of life and relationships. Thus, there is this pressure to bridge the gap between the frequency of the sexual difficulties and the minority that actually goes for professional help. This shows that there are really serious and basic issues in the way we perceive, diagnose, and provide accessible and comprehensive treatment for sexual health concerns.

Having clear and consistent ways to classify sexual dysfunctions is essential for research, setting diagnostic standards, and creating effective treatment plans. Each classification requires a simple definition of a sexual difficulty. The sexual dysfunctions are mainly categorised by the sexual difficulty that occurs along the four stages of the sexual response cycle, i.e., desire, arousal, orgasm, and resolution, and also conditions concerned with sexual pain. A person could have lived with the issue for

life, or it comes about later in life from a certain period where everything was functioning normally. It could be situational or dependent on certain circumstances or specific partners, or it could be more generalized. Being able to understand the dysfunction we are dealing with calls for understanding the sexual dysfunction in the context of the relationships they are associated with. An absence of sexual response in some cases could be a rational or protective reaction to certain difficult situations (such as relationship conflicts or high-stress levels) rather than being her or his personal failure. Hamzat et al. (2025) noted that expressed emotions towards adolescents could affect how they handle their sexuality and other related dysfunction in their lives

This paper aims to explore sexual disorders and dysfunction thoroughly, moving beyond just looking at the symptoms. We aim to examine the many factors that contribute to these issues. Our main goals include defining and categorising common sexual dysfunctions, carefully examining the different pathways that lead to them, and examining established theories that help explain how they develop and persist. Our approach is primarily based on a review of existing academic research. This integrated view will consistently highlight how biological factors, psychological states, and social and cultural influences all work together to shape a person's sexual health. The main argument of this paper is that to understand and effectively manage sexual disorders truly, we need a holistic perspective one that smoothly combines biological, psychological, and social aspects, moving beyond simple explanations to embrace the full, rich complexity of human sexuality.

THEORETICAL AND CONCEPTUAL FRAMEWORK

To truly make sense of sexual disorders and dysfunction, we need a solid theoretical foundation that recognises their many different roots.

1. *Psychodynamic Theory*: particularly influenced by the ground-breaking work of Sigmund Freud, offers deep insights into the often-unconscious processes and early life experiences that might contribute to sexual dysfunction. Freud's idea of psychosexual development, moving through oral, anal, phallic, latency, and genital stages, suggests that unresolved conflicts

Or lingering issues from any of these earlier life phases can show up as sexual challenges in adulthood. For example, difficulties in successfully navigating the oedipal complex or the lasting effects of early childhood trauma might unconsciously block healthy sexual expression or lead to specific dysfunctions (Freud, 1962). While we acknowledge that psychodynamic ideas can be tough to test directly through experiments, they powerfully highlight the profound, often hidden, impact of past experiences and unconscious motivations on a person's current sexual functioning.

2. *Cognitive-Behavioural Theories (CBT)*: provides an incredibly useful way to understand how a person's thoughts, feelings, and actions work together in a continuous loop to influence sexual function. The cognitive part of this theory focuses on unhelpful beliefs, persistent negative thoughts, and paralyzing performance anxiety.

All of these can unfortunately create a self-fulfilling prophecy of sexual failure (Beck, 1976). For instance, if someone has catastrophic thoughts about their sexual performance, it can quickly make their anxiety worse, which then directly prevents physical arousal. The behavioral side of CBT looks at learned patterns, like avoiding sexual situations or following unhelpful, ingrained "scripts" for sex. As a result, CBT

interventions typically involve actively challenging and changing irrational thoughts, developing new and healthier ways to cope, and doing structured behavioral exercises designed to help people re-establish healthy and satisfying sexual patterns gradually.

Finally, Human Sexual Response Models offer a crucial physiological and psychological roadmap for understanding the step-by-step progression of sexual activity.

3. *The Biopsychosocial Model*: is a core, widely accepted framework that suggests our sexual health, and conversely, any dysfunctions, come from a dynamic interaction between biological, psychological, and social factors (Engel, 1977). On the biological side, this includes things like intricate brain functions, shifting hormone levels, the health of our blood vessels, and even the often-overlooked effects of various medications or long-term illnesses. Psychologically, it covers a person's unique life experiences, their deep-seated beliefs about sex, their current emotional state (like feeling anxious or depressed), past traumas, and the subtle but powerful dynamics within their relationships. Socially, cultural norms, strong religious beliefs, how someone was raised, and broader societal pressures all significantly shape sexual attitudes and behaviors, often leading to distress or dysfunction. This comprehensive model strongly emphasizes that any truly effective assessment and treatment plan must carefully consider all three of these interconnected areas.

4. *The traditional four-phase model*: as proposed by Masters and Johnson (1966), represents a linear succession: Desire (the initial urges of sexual connection, inclusive of thoughts or fantasies), Arousal/Excitement (when blood is engorged in the genitalia: a female's genitals are lubricated, and a male has an erection), Orgasm (characterized by involuntary rhythmic contractions of the

skeletal muscle with release of tension), and Resolution (being the body's return to an unaroused state). However, with research over the years, especially from Helen Singer Kaplan's work (1979), it is suggested that the sexual responses of women are often more intricate and non-linear.

For women, sexual response might involve phases that overlap and happen in a varied order, a pattern especially common in long-term relationships. Also, desire can often be reactive instead of purely spontaneous, signifying that desire might emerge and grow during sexual activities themselves, often propelled by motives that may have nothing to do with sex: to feel close to a significant other, to please him, or even as a way of unwinding (Basson, 2005; Carvalheira et al., 2010) Further, an important contrast between men and women is that for women, physical signs of arousal (such as genital lubrication) might not automatically correspond to their feelings of sexual excitement; in women, thoughts and emotions can have a profound effect on their subjective feelings of sexual excitement much more so than just the physical signs alone.(Birnbaum, n.d.; Brotto et al., 2010).

These foundational models remain essential for accurately classifying and diagnosing specific sexual dysfunctions based on which part of the sexual response cycle is affected.

Definition and Diagnostic Categorization

Sexual disorders, also known as sexual dysfunctions, are conditions that prevent individuals or couples from achieving satisfaction during coitus. They may be lifelong or acquired, generalised or situational, and due to biological, psychological, or social factors. The most widely utilized classification systems in clinical and research settings are the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), and the

International Classification of Diseases, Eleventh Revision (ICD-11).

Now, according to DSM-5 and ICD-11, the DSM-5 gives specific diagnostic features for sexual dysfunctions. These include the presence of symptoms for a minimum of six months, significant distress caused by the symptoms, and that the dysfunction is not explainable by a non-sexual mental disorder, extreme relationship distress, or other stressors. Sexual dysfunctions are also categorized under a distinct chapter in the ICD-11 and emphasize the duration, distress, and lack of other medical causes. The major categories include;

1. Sexual Desire Disorders:

Hypoactive sexual desire disorder (HSDD): characterized by a persistent or recurrent deficiency in sexual thoughts and interest. It is more commonly diagnosed in females but can also occur in males. Low desire is normal as a function of age, but is not a disorder unless it causes distress.

2. Arousal Disorders:

Female Sexual Arousal Disorder: Marked by an inability to attain or maintain sufficient lubrication and genital swelling upon the onset or continuation of sexual excitement or activity.

Erectile Disorder: Marked by a repeated inability to attain or maintain an erection sufficient for satisfactory sexual performance. It is more common in older males and frequently coexists with other medical conditions.

3. Orgasmic Disorders

Delayed Ejaculation: Markedly delayed or inability to ejaculate in men despite proper stimulation and desire.

Female Orgasmic Disorder: Difficulty or inability to have orgasm during sexual

activity. A variety of psychological and interpersonal factors influences it.

4. Sexual Pain Disorders

- *Genito-Pelvic Pain/Penetration Disorder*: Comprises vaginal penetration issues, pelvic pain, fear of or anxiety about pain, and tensing of the muscles of the pelvic floor. It occurs only in women and exists across different age groups.

5. Paraphiliac Disorders:

- *Exhibitionistic Disorder*: Is a disorder that constitutes intense sexual arousal by exposing one's genitals to an unsuspecting person.

- *Pedophilic Disorder*: It is characterised by recurrent sexual urges, fantasies, or behaviours toward prepubescent children. Diagnosis requires that the individual be at least 16 years old and at least 5 years older than the child.

Discussion

Gender Differences and Age-Related Prevalence:

Sexual disorders manifest differently across genders. Desire and arousal disorders are reported more often by women, while men preferentially report erectile and ejaculation disorders. Age has a general effect on prevalence, as older individuals are more likely to develop arousal-type dysfunctions secondary to physiological changes.

Causes of Sexual Disorders

The aetiology of sexual disorders is multifactorial and comprises biological, psychological, social, and substance-related factors. They occur independently or interplay and produce various kinds of sexual dysfunction. To diagnose accurately,

treat effectively, and reduce stigma, it is necessary to understand causation.

A. Biological Causes: Biological etiology of sexual dysfunction typically encompasses hormonal imbalance, neurological disorders, vascular disease, and chronic illness. Hormones such as testosterone, estrogen, and prolactin have a role to play in regulating sexual desire and function. Hypogonadism, for instance, will lead to decreased libido in both sexes. Neurological diseases like multiple sclerosis, Parkinson's disease, or spinal cord injury can disrupt nerve transmission critical to sexual response. Additionally, vascular disease, such as atherosclerosis or diabetes-related endothelial dysfunction, may impair blood flow, contributing to erectile or arousal dysfunction. Chronic illnesses such as cancer, renal failure, or chronic pain syndromes are also associated with compromised sexual function due to fatigue, pain, or treatment side effects.

B. Psychological Causes: Psychological factors are common and often the primary aetiology of sexual disorders. Illnesses such as anxiety, depression, and a history can significantly influence sexual function. Anxiety, particularly performance anxiety, can interfere with arousal and orgasm. Depression is most commonly expressed in losing interest in activities one used to enjoy, such as sexual ones, and can be exacerbated by antidepressant medication. Trauma or sexual abuse histories can lead to avoidance symptoms, intimacy phobia, and hypervigilance during sex. Cognitive distortions, a negative view of the body, and low self-esteem make the emotional backdrop of sexual dysfunction difficult.

C. Social and Cultural Factors: Social and cultural variables are primarily responsible for shaping sexual attitudes and behavior. Relationship problems that could lead to desire and arousal disorders, such as poor communication, conflict of issues, or infidelity, are among these factors. Cultural

or religious sex attitudes that stigmatise sex can produce guilt or shame that inhibit sexual expression. Inhibiting sexuality is typical in societies with rigid gender roles or sexuality taboos, where individuals conceal their sexual feelings or experience identity-based issues. Abuse, emotional, physical, or sexual, within or outside the context of a close relationship, can also hurt sexual functioning. Several young women hide under being a single/ baby mama to hide their psychological dysfunction (Akinwale et al 2025).

D. Substance-Induced Sexual Disorders:

Drugs, alcohol, and certain medications can induce or exacerbate sexual dysfunctions. Chronic alcohol use is linked with reduced libido and erectile dysfunction. Use of drugs for recreational purposes, e.g., cocaine, opioids, and cannabis, can disrupt arousal and orgasmic function. Medications such as selective serotonin reuptake inhibitors (SSRIs), antihypertensive drugs, and antipsychotics have been found to cause sexual side effects. Substance-induced sexual disorders are most commonly reversible upon cessation of the substance. Still, long-term use may create irreversible effects or co-occur with psychiatric illness that must be treated at the same time.

Assessment and Diagnosis

Diagnosis and evaluation of sexual disorders must be multi-disciplinary, sensitive, and extensive. Since sexual health issues are private and often stigmatized, clinicians must establish a non-judgmental and trustworthy environment to acquire accurate information.

A thorough sexual history is important in the assessment of the nature, extent, and context of the sexual dysfunction. This involves the following questions about the onset of symptoms, relationship, emotional relationship with partners, prior sexual experiences, and current sexual activity.

The cultural values and individual beliefs concerning sexuality should also be obtained during the history taking.

Standardised questionnaires provide clinicians with unambiguous data to assist in diagnosis and also in progress monitoring. Among the questionnaires are the Female Sexual Function Index (FSFI) and International Index of Erectile Function (IIEF), which assess various aspects of desire, arousal, lubrication or erection, orgasm, satisfaction, and pain. The questionnaires help to identify specific dysfunctions and are applied in clinical and research settings.

There are some medical tests required to rule out physiological etiologies for sexual dysfunction. These may involve hormonal evaluation, vascular assessment, neurological testing, and review of current medications. Additionally, psychological testing determines the underlying issues of depression, anxiety, trauma, or relationship stressors that may be responsible for the dysfunction.

The evaluation should be conducted with sensitivity to culture. Clinicians must know how sexual attitudes, sexual behaviour, and acceptance of talking about sexual concerns are affected by religious and cultural beliefs. Hesitancy to report concerns will be encountered in conservative patients, and some cultures may normalize clinically significant symptoms.

Even with exposure to clinical guidelines and resources, numerous challenges remain. Patients may feel shy or unwilling to be honest, and therefore underreport. Clinicians themselves may not receive adequate training or time to conduct appropriate sexual screening. Thus, improving clinician education and integrating sexual health into routine assessments represent essential moves toward improved diagnosis and care.

Treatment and Management Approaches

Treatment and care of sexual disorders are an issue of individualised and often multidisciplinary care, combining medical, psychological, and relational treatments. Treatment ought to be aimed at the original cause and consider the patient's choice, cultural background, and relationship.

A. Psychotherapeutic Approaches

Psychotherapeutic treatment remains the mainstay of the therapy of sexual disorders, especially the psychological or relational ones. Cognitive Behavioral Therapy (CBT) helps the patient to become aware of and modify skewed thinking, beliefs, and behaviors that affect sexual performance and gratification. It can cure performance anxiety, for example, negative body image, or guilt over sex.

Sex therapy, typically conducted by trained sex therapists, employs techniques like sensate focus a series of exercises under guidance, which reduce performance anxiety and rebuild intimacy. It is particularly beneficial for couples with desire differences, arousal disorders, or orgasmic disorders.

Couples therapy addresses the relational issues of sexual dysfunction. It focuses on improving communication, conflict resolution, and emotional intimacy, which can enhance sexual satisfaction.

B. Pharmacological Interventions

Pharmacotherapy works for specific types of sexual disorders. Phosphodiesterase type 5 (PDE5) inhibitors such as sildenafil (Viagra), tadalafil (Cialis), and vardenafil (Levitra) are the first-line treatment for erectile dysfunction. They cause improved blood flow to the penis that helps in erection during sexual stimulation.

Pharmacologic treatments can be administered for hormonal imbalances. Testosterone replacement therapy in males with hypogonadism and female decreased libido, and estrogen therapy for the treatment of vaginal dryness and pain in postmenopausal women are a few examples. Side effects of pharmacologic therapy are headache, flushing, dizziness, and infrequently, more serious cardiovascular complications. Adequate education of the patient about the use and monitoring for complications is required.

C. Medical and Physical Treatments

Physical treatments are helpful for certain patients. Vacuum erection devices create a suction that helps penile engorgement, often used by men who cannot use oral therapy. Surgical interventions include penile implants for severe erectile dysfunction and reconstructive surgery for anatomical deformity.

Pelvic floor physical therapy also suits patients, especially women, who have genito-pelvic pain/penetration disorder. They strengthen pelvic muscles, reduce pain, and improve sexual functioning.

D. Integrated/Multidisciplinary Approaches

The most effective method is one that incorporates medical, psychological, and social treatments. Consultations among urologists, gynecologists, endocrinologists, psychologists, and sex therapists are coordinated in such a manner that all the etiologies are addressed.

This interdisciplinary approach increases the efficacy of treatment as well as patient satisfaction. Nonetheless, compliance may be affected by barriers including access limitations to specialists, expense, and unawareness. Enhancing accessibility through telemedicine and programs at the

community level may enhance participation.

In conclusion, sexual disorder management is necessitated by a comprehensive and individualised program integrating patient-centred care and evidence-based treatment. Multidisciplinary management not only addresses the complexity of sexual dysfunctions but also promotes long-term outcomes and overall wellness

Cultural and Gender Variations

Sexuality is deeply rooted within culture and societal norms that massively affect the perception, expression, and diagnosis of sexual disorders. Cultural perceptions determine what is regarded as deviant or accepted sexual behavior and go a long way in deciding whether people seek help or tell symptoms of dysfunction to another person (Bhugra & De Silva, 2016). In the very conservative societies, sexual disorders are thought to be moral lapses or spirit sickness, the kinds of things that add to stigma and silence (Okulate & Olayinka, 2021). This sinking of culture pours into the deep wells of underreporting and disintermediation for diagnosis and treatment.

Interpretation and taboos on sexual health topics also act as impediments in clinical assessments: it is said that even in very medical settings, many patients do not feel at ease or develop a sense of shame, with women staying more prone to this due to circumstances that weigh on sexual behavior and modesty-gendered in certain perspectives (Gott & Hinchliff, 2003). In patriarchal societies, female sexual dysfunction could somehow be discounted or disregarded, whereas male dysfunction is medically pathologised owing to social imperatives surrounding masculinity and performance (Wylie & Barrett, 2007).

Sexual dysfunctions may present differently in each gender, as they depend on biological and psychosocial factors. Men

usually complain of erectile dysfunction or delayed ejaculation, while women often complain of disorders of arousal, low sexual desire, and pain problems (Clayton et al., 2017). Gender norms set differential levels of acceptability or help-seeking behavior in the opposite sex.

Sexual health care poses difficult problems for LGBTQ+ persons. Non-heteronormative sexual identities are often seen as reputable pathologies or misunderstood by providers without a cultural competence, which leads to misdiagnosis or overt neglect (Pachankis & Jackson, 2021). Transgender persons may, in particular, experience sexual dysfunction induced by testosterone or estrogen treatments, by surgical procedures, or simply by body dysphoria, good-sexual-dysfunction-care-affirming-starved (Wierckx et al., 2011).

To combat these disparities, assessments and interventions must be capable of recognizing and respecting diversities of sexual identity, sexual behaviour, and sexuality norms. Clinicians should be sensitized to the intersection of gender, culture, and sexuality to provide a diagnosis and treatment which is inclusive.

Impact on Quality of Life

Sexual dysfunctions exert a profound impact on emotional and psychological well-being, often triggering feelings of frustration, shame, inadequacy, and depression (Laumann et al., 1999). Such emotional responses may increase the disorder itself and may generate a vicious circle of anxiety-retreat (Rosen & Bachmann, 2008). Sexual dysfunctions can cause psychological trauma, isolation, and even suicidal thoughts when coupled with other mental health issues.

Adding to common distressing issues is strained relationships and marriages, since sexual intimacy is an essential part of many

romantic relationships. The dysfunction may interfere with tenderness between the partners, disrupt communication, and ultimately cause the partners to drift from each other (McCabe & Connaughton, 2014). Furthermore, a severe, unresolved dysfunction may give room to infidelity or even the breakup of the relationship.

Sexual problems also affect self-esteem and identity, especially when sexuality is central to one's self-concept or cultural expectations regarding masculinity/femininity. The longer the dysfunction lasts, the more confidence a person loses and develops a bad self-image to increase sexual anxiety (Bancroft, 2009).

The significance of sex disorders for work and one's social life is often underestimated. A person might be distracted or fatigued or the emotional manifestation of an interpersonal stressor may reduce motivation enough to impair job performance or increase absenteeism (Derogatis & Meyer, 1979). Also, romantic or social relationships may suffer as individuals begin withdrawing from social activities because of embarrassment or fear of being exposed.

Thus, because of the wide range of effects, sexual dysfunction must be considered not just a clinical issue but also an important quality-of-life issue. A holistic approach to treating sexual dysfunction is thus required, one in which relational, psychological, and lifestyle approaches are intertwined.

Current Trends and Research Developments

New therapies and fresh hope for the sexually-disordered; sexual health is rapidly advancing. Mindfulness-based therapy has shown to be a workaround for sexual desire, arousal, and satisfaction issues in women, especially those who have undergone trauma (Brotto et al., 2012). Neuromodulation therapies have also been

proposed to rectify cases of psychogenic erectile dysfunction and other sexual dysfunctions that arise from neural dysregulations (Zhao et al., 2021).

Digital health tools like mobile apps, virtual therapy, and telemedicine platforms are now being used to provide sexual therapy and education in a discrete and easy-to-access manner (Anderson et al., 2020). Such technologies favor those living in rural or conservative settings, where face-to-face interaction is scarce or frowned upon. Akinwale et al (2025) in a study on psychosocial intervention noted that psychological intervention is a critical domain for overall health.

Changing societal attitudes bring with them changes in education for sexuality. Comprehensive sex education that is inclusive in schools and community programs has been associated with decreased sexual dysfunction, earlier help-seeking, and better relationship outcomes (Kirby, 2008). There is more and more advocacy for the inclusion of sexual health in a general wellness framework to foster a sex-positive and non-judgmental discourse.

Along with this, the most recent research explores the intersectional nature of sexuality-that is, how age, disability, race, and sexual orientation sites affect sexual function and intersect with access to care (Meston et al., 2020). This has triggered a turn toward models of treatment that are culturally informed and individualized.

As the research in this area continues to grow, the integration of technology, psychology, and social awareness continues to reshape sexual health care, with the result being more enhanced, empathetic, and inclusive practices.

Conclusion

Sexual disorders and dysfunctions are complicated and deeply personal issues

affecting a vast spectrum of persons, differing in terms of gender, age, or culture. As has also been explored in this paper, such disorders can be caused by a multitude of reasons, namely biological, psychological, relational, social, or cultural. More often than not, these factors intermingle with each other in complex relations that present great challenges in treatment and diagnosis. Such disorders interfere with sexual satisfaction, interfere with the fostering of self-esteem in client relationships, as well as with clients' general emotional states.

To understand sexual dysfunction, one must take on a full biopsychosocial approach that includes all aspects of the person's life. Beliefs, history, relations, and physiology should all be taken into account. Consequently, diagnosis requires more than just an evaluation of the symptoms; rather, it should involve a culturally sensitive approach along with using adequately validated instruments, such as the FSFI and IIEF. In contrast, treatment has to be targeted toward the individual and be a combination of psychotherapeutic, medical, and usually some form of relationship- or sex-oriented therapies.

Stigma and cultural taboos have imposed great restrictions against open discussion and seeking timely help. The establishment of sexual health as an integral part of general health, thus an area that must be destigmatised, sets an immediate demand for sexual health training in medical settings.

The ultimate redemption of sexual disorders is no longer just a clinical matter; it is the restoration of the human spirit, restoring confidence in partnerships and therefore goes one step further into building an inclusive, healthy perception of human sexuality itself. Upon creating awareness, reducing stigma, and embracing comprehensive care, we become a step closer to having sexual health identified, respected, and supported as an essential part of life.

REFERENCE

- Akinwale, G. A., & Oloko, T. O. (2025). Baby mama syndrome: Psychosocial and developmental implications of an emerging trend. *Acta Universitatis Sapientiae, Social Analysis*, 15, 140–156. <https://doi.org/10.47745/aussoc-2025-0008>
- Akinwale, G. A. & Ojakorotu, Victor (2025). Human at the Centre Stage of Peacebuilding: A review of the Use of Psychosocial Intervention in Africa. *Journal of. Biomedical. Research. Vol. 28(4s) (November 2025); 549-554 Research Article in Lagos.*
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.).
- Anderson, M., McKee, M., & Nelson, J. (2020). Digital health interventions to improve sexual health: Scoping review. *Journal of Medical Internet Research*, 22(6), e16724. <https://doi.org/10.2196/16724>
- Auctores Online. (2024). Sexual disorders in ICD-11: Innovations and their discussion. Auctores Online. <https://auctoresonline.org/article/sexual-disorders-in-icd-11-innovations-and-their-discussion>
- Bancroft, J. (2009). Human sexuality and its problems (3rd ed.). Elsevier Health Sciences.
- Basson, R. (2005). Women's sexual dysfunction: revised and expanded definitions. *Canadian Medical Association Journal*, 172, 1327-1333.
- Beck, A. T. (1976). Cognitive therapy and the emotional disorders. International Universities Press.
- Bhugra, D., & De Silva, P. (2016). Sexual dysfunction across cultures. In P. De Silva & S. Rachman (Eds.), *Abnormal and clinical psychology: An introductory textbook* (pp. 322–336). Routledge.
- Birnbaum, G. E. (n.d.). Sexual Dysfunctions. Interdisciplinary Center (IDC) Herzliya.
- Brotto, L. A., Basson, R., & Luria, M. (2012). A mindfulness-based group psychoeducational intervention targeting sexual arousal disorder in women. *Journal of Sexual Medicine*, 9(6), 1444–1458. <https://doi.org/10.1111/j.1743-6109.2012.02703.x>
- Brotto, L. A., Bitzer, J., Laan, E., Leiblum, S., & Luria, M. (2010). Women's sexual desire and arousal disorders. *Journal of Sexual Medicine*, 7(1 Pt 2), 586-614. (Published correction appears in J Sex Med. 2010;7(2 Pt 1):856).
- Carvalho, A. A., Brotto, L. A., & Leal, I. (2010). Women's motivations for sex: exploring the diagnostic and statistical manual, fourth edition, text revision criteria for hypoactive sexual desire and female sexual arousal disorders. *Journal of Sexual Medicine*, 7(4 Pt 1), 1454-1463.
- Clayton, A. H., & Balon, R. (2009). The impact of mental illness and psychotropic medications on sexual functioning: The evidence and management. *The Journal of Sexual Medicine*, 6(5), 1200–1211. <https://doi.org/10.1111/j.1743-6109.2009.01215.x>
- Clayton, A. H., Kingsberg, S. A., & Goldstein, I. (2017). Evaluation and management of female sexual dysfunction. *Mayo Clinic Proceedings*, 92(4), 562–578. <https://doi.org/10.1016/j.mayocp.2016.11.002>

- Derogatis, L. R., & Meyer, J. K. (1979). Psychological factors in impotence. *Psychiatric Clinics of North America*, 2(2), 319–332.
- Engel, G. L. (1977). The need for a new medical model: A challenge for biomedicine. *Science*, 196(4286), 129-136.
- Freud, S. (1962). Three essays on the theory of sexuality (J. Strachey, Trans.). Basic Books. (Original work published 1905).
- Gott, M., & Hinchliff, S. (2003). Barriers to seeking treatment for sexual problems in primary care: A qualitative study with older people. *Family Practice*, 20(6), 690–695. <https://doi.org/10.1093/fampra/cm612>
- Hamzat, M.F., Aroyewun, B.A. & Akinwale, G. (2025). Development and validation of the perceived expressed emotion scale for adolescents (PEEs-GAP). *BMC Psychiatry* 25, 268 <https://doi.org/10.1186/s12888-025-06695-6>
- Kaplan, H. S. (1979). Disorders of sexual desire and other new concepts of sex therapy. Brunner/Mazel.
- Kirby, D. (2008). The impact of sex and HIV education programs in schools and communities on sexual behaviors among young adults. *Sexuality Research and Social Policy*, 5(3), 24–40. <https://doi.org/10.1525/srsp.2008.5.3.24>
- Laumann, E. O., Paik, A., & Rosen, R. C. (1999). Sexual dysfunction in the United States: Prevalence and predictors. *JAMA*, 281(6), 537–544. <https://doi.org/10.1001/jama.281.6.537>
- Levine, S. B. (2003). The nature of sexual desire: A clinician’s perspective. *Archives of Sexual Behavior*, 32(3), 279–285. <https://doi.org/10.1023/A:1023427823704>
- Masters, W. H., & Johnson, V. E. (1966). Human sexual response. Little, Brown.
- McCabe, M. P., & Connaughton, C. (2014). Psychosocial factors in sexual dysfunction among men and women. *Current Psychiatry Reports*, 16(11), 443. <https://doi.org/10.1007/s11920-014-0443-0>
- Meston, C. M., Chivers, M. L., & Rellini, A. (2020). Women’s sexual dysfunctions: Progress in diagnosis and treatment. *Annual Review of Clinical Psychology*, 16, 233–256. <https://doi.org/10.1146/annurev-clinpsy-050718-095538>
- Okulate, G. T., & Olayinka, M. O. (2021). Culture and sexual dysfunction: A review of the Nigerian perspective. *Nigerian Journal of Psychiatry*, 19(1), 10–17.
- Pachankis, J. E., & Jackson, S. D. (2021). Sexual orientation and mental health: Advances in research and evidence-based intervention. *Annual Review of Clinical Psychology*, 17, 481–505. <https://doi.org/10.1146/annurev-clinpsy-081219-105748>
- Reed, G. M., Drescher, J., Krueger, R. B., Atalla, E., Cochran, S. D., First, M. B., ... & Saxena, S. (2016). Disorders related to sexuality and gender identity in the ICD-11: Revising the ICD-10 classification based on current scientific evidence, best clinical practices, and human rights considerations. *World Psychiatry*, 15(3), 205–221. <https://doi.org/10.1002/wps.20354>

- Rosen, R., Brown, C., Heiman, J., Leiblum, S., Meston, C., Shabsigh, R., ... & Ferguson, D. (2000). The Female Sexual Function Index (FSFI): A multidimensional self-report instrument for the assessment of female sexual function. *Journal of Sex & Marital Therapy*, 26(2), 191–208. <https://doi.org/10.1080/009262300278597>
- Rosen, R. C., Riley, A., Wagner, G., Osterloh, I. H., Kirkpatrick, J., & Mishra, A. (1997). The International Index of Erectile Function (IIEF): A multidimensional scale for assessment of erectile dysfunction. *Urology*, 49(6), 822–830. [https://doi.org/10.1016/S0090-4295\(97\)00238-0](https://doi.org/10.1016/S0090-4295(97)00238-0)
- Wierckx, K., Van Caenegem, E., Elaut, E., Van Hoorde, B., Heylens, G., De Cuypere, G., & T'Sjoen, G. (2011). Sexual desire in trans persons: Associations with sex reassignment treatment. *Journal of Sexual Medicine*, 8(1), 337–345. <https://doi.org/10.1111/j.1743-6109.2010.01957.x>
- World Health Organization. (2019). International classification of diseases for mortality and morbidity statistics (11th ed.). <https://icd.who.int/>
- Wylie, K. R., & Barrett, J. (2007). The psychosocial impact of erectile dysfunction. *British Journal of Nursing*, 16(3), 148–152. <https://doi.org/10.12968/bjon.2007.16.3.22951>
- Zhao, Z., Zhang, L., Wang, H., & Tang, Z. (2021). Transcranial magnetic stimulation for sexual dysfunction: A systematic review. *Frontiers in Neuroscience*, 15, 694763. <https://doi.org/10.3389/fnins.2021.694763>